

Dissertation Training

at

International Institute of Health Management Research New Delhi

Birth preparedness and complication readiness among pregnant and recently delivered women in India: Systematic review and Meta-analysis.

By

Miss. Tanya Singh

PG/21/120

Under the guidance of

Dr. Sidharth Sekhar Mishra

PGDM (Hospital & Health Management)

2021-23



**International Institute of Health Management Research
New Delhi**

Dissertation Training

at

International Institute of Health Management Research New Delhi

Birth preparedness and complication readiness among pregnant and recently delivered women in India: Systematic review and Meta-analysis.

By

Miss. Tanya Singh

PG/21/120

Under the guidance of

Dr Sidharth Sekhar Mishra

PGDM (Hospital & Health Management)

2021-23



**International Institute of Health Management Research
New Delhi**

The certificate is awarded to

Miss. Tanya Singh

in recognition of having successfully completed

her Dissertation

at IIHMR, Delhi

and has successfully completed her Project on

Birth preparedness and complication readiness among pregnant and recently delivered women in India: Systematic review and Meta-analysis.

Date- 15th January to 15th April, 2023

International Institute of Health Management Research New Delhi

She comes across as a committed, sincere & diligent person who has

a strong drive & zeal for learning.

We wish her all the best for future endeavours.

Training & Development

Zonal Head-Human Resources

TO WHOMSOEVER IT MAY CONCERN

This is to certify that **Miss. Tanya Singh**, student of PGDM (Hospital & Health Management) from International Institute of Health Management Research, New Delhi has undergone internship training at **International Institute of Health Management Research New Delhi** from **15th January to 15th April, 2023**.

The Candidate has successfully carried out the study designated to her during internship training and her approach to the study has been sincere, scientific and analytical.

The Internship is in fulfilment of the course requirements.

I wish her all success in all her future endeavours.

Dr. Sumesh Kumar
Associate Dean, Academic and Student Affairs
IIHMR, New Delhi


Dr. Sidharth Sekhar Mishra
Assistant Professor
IIHMR, New Delhi

Certificate of Approval

The following dissertation titled “**Birth Preparedness and Complication Readiness among Pregnant Women and Recently Delivered Women in India: Systematic Review and Meta-Analysis**” is hereby approved as a certified study in management carried out and presented in a manner satisfactorily to warrant its acceptance as a prerequisite for the award of **PGDM (Hospital & Health Management)** for which it has been submitted. It is understood that by this approval the undersigned do not necessarily endorse or approve any statement made, opinion expressed or conclusion drawn therein but approve the dissertation only for the purpose it is submitted.

Dissertation Examination Committee
for evaluation of dissertation.

Name

Signature

Sidharth Sekhar
Mishra

Sidharth Sekhar Mishra

Rupla Banerjee

Rupla

J. P. Mishra

J.P.

Certificate from Dissertation Advisory Committee

This is to certify that **Miss.Tanya Singh**, a graduate student of the **PGDM (Hospital & Health Management)** has worked under our guidance and supervision. She is submitting this dissertation titled **“BIRTH PREPAREDNESS AND COMPLICATION READINESS AMONG PREGNANT AND RECENTLY DELIVERED WOMEN IN INDIA: SYSTEMATIC REVIEW AND META-ANALYSIS”** in partial fulfilment of the requirements for the award of the **PGDM (Hospital & Health Management)**.

This dissertation has the requisite standard and to the best of our knowledge no part of it has been reproduced from any other dissertation, monograph, report or book.

Sidharth Sekhar Mishra


Dr Sidharth Sekhar Mishra,
Assistant Professor,
IIHMR, New Delhi

Sidharth Sekhar Mishra

Dr Sidharth Sekhar Mishra
Assistant Professor,
IIHMR, New Delhi

CERTIFICATE BY SCHOLAR

This is to certify that the dissertation titled “**Birth preparedness and complication readiness among pregnant and recently delivered women in India**”: **Systematic review and Meta-analysis**” and submitted by **Miss.Tanya Singh**, Enrollment No. **PG/21/120** under the supervision of **Dr Sidharth Sekhar Mishra**, Assistant Professor, IIHMR Delhi for award of **PGDM** (Hospital & Health Management) of the Institute carried out during the period from **15th January to 15th April, 2023** embodies my original work and has not formed the basis for the award of any degree, diploma associate ship, fellowship, titles in this or any other Institute or other similar institution of higher learning.



Signature

FEEDBACK FORM

Name of the Student: Miss. Tanya Singh

Name of the Organisation in Which Dissertation Has Been Completed: International Institute of Health Management Research New Delhi

Area of Dissertation: Birth preparedness and complication readiness among pregnant and recently delivered women in India: Systematic review and Meta-analysis

Attendance: Perfect adherence to dissertation norms

Objectives achieved: The student understood the details of the Systematic Review and Meta-Analysis

Deliverables: As per dissertation guidelines

Strengths: Sincerity, attention span, concentration, hard work, diligence, detail-orientedness

Suggestions for Improvement: Wait for the question to end before

Suggestions for Institute (course curriculum, industry interaction, placement, alumni):

PLEASE DEVELOP A+ BATCH FOR SUCH GOOD STUDENTS.

Sidhant Keshav Mishra

Signature of the Officer-in-Charge/ Organisation Mentor (Dissertation)

Date: 28th June, 2023

Place: I I H M R, Delhi



INTERNATIONAL INSTITUTE OF HEALTH MANAGEMENT RESEARCH (IIHMR)

Plot No. 3, Sector 18A, Phase- II, Dwarka, New Delhi- 110075

Ph. +91-11-30418900, www.iihmrdelhi.edu.in

CERTIFICATE ON PLAGIARISM CHECK

| | | | |
|--|--|-------------------|-----------|
| Name of Student (in block letter) | Ms. TANYA SINGH | | |
| Enrolment/Roll No. | PG/21/120 | Batch Year | 2021-2023 |
| Course Specialization (Choose one) | | Health Management | |
| Name of Guide/Supervisor | Dr. Sidharth Sekhar Mishra | | |
| Title of the Dissertation/Summer Assignment | Birth preparedness and complication readiness among pregnant and recently delivered women in India: Systematic review and Meta-analysis. | | |
| Plagiarism detects software used | "TURNITIN" | | |
| Similar contents acceptable (%) | Up to 15 Percent as per policy | | |
| Total words and % of similar contents Identified | 11% | | |
| Date of validation (DD/MM/YYYY) | 28 June 2023 | | |

Guide/Supervisor

Sidharth Sekhar Mishra

Name: Dr. Sidharth Sekhar Mishra

Signature:

Report checked by

Institute Librarian

Signature:

Date:

Library Seal

Ashok Kumar



Student

Name: Ms. Tanya Singh

Signature:

Tanya Singh

Dean (Academics and Student Affairs)

Signature:

Date:

(Seal)

ACKNOWLEDGEMENT

It is an esteemed pleasure to present this research project by thanking each and everyone who helped me in this task. I would like to express my sincere gratitude towards my guide **Dr. Sidharth Sekhar Mishra**, Assistant Professor, IIHMR Delhi, who helped me immensely throughout the tenure of my internship. He inspired me greatly to work in this project with his valuable guidance, support, interest, encouragement, involvement and advice.

I would like to thank Dr Sidharth Sekhar Mishra, Assistant Professor, IIHMR Delhi, for allowing me to experience such great opportunities and for providing data for my learning.

I would also like to express my special thanks to **Dr. Sutapa Neogi, (Director)** IIHMR, Delhi and **Dr. Anuj Kumar Pandey** and **Dr. Diksha Gautam** for their support and guidance in providing such great opportunity which helps in to grow and learn about many interesting aspects.

List of Table and Figures-

| | |
|--|----|
| Figure 1:- Conceptual Diagram of How BPCR may increase the use of skilled care | 19 |
| Figure 2:- PRISMA Flow chart summarizing the article selection process | 30 |
| Figure 3:- Forest Plot Depicting Overall BPCR status in India | 32 |
| Figure 4 :-Forest plot Depicting (Weight of Studies). | 34 |
| Figure 5: -Forest plot Depicting subgroup analysis – Delivery Status wise | 36 |
| Figure 6:- Forest plot Depicting subgroup analysis – Region wise..... | 37 |
| Figure 7:- Funnel Plot..... | 40 |
| Figure 8:- Funnel Plot after Performing Eggers Test..... | 40 |
| Figure 9:- Stacked bar chart depicting ROB. | 41 |
| Table 1:-PICO Format..... | 24 |
| Table 2:- Prevalence of components of BP and CR..... | 39 |

| So. No. | <u>Table of Contents</u> | Page No. |
|--------------------|--|---------------------|
| | Introduction..... | 17 |
| | Rationale | 21 |
| | Methodology..... | 23 |
| | ... | 23 |
| | Study Registration | 23 |
| | Search strategy | 24 |
| | PICO Format | 24 |
| | Study Selection | 24 |
| | Selection Process | 24 |
| | Data extraction and data Collection | 25 |
| | Main Outcome | 25 |
| | Measure of effect | 26 |
| | Risk of Bias (ROB) Quality Assessment | 26 |
| | Data synthesis, statistical analysis, and investigation of | 27 |
| | Heterogeneity | 27 |
| | Sub-group analysis | 27 |
| | Definition of BPCR | 28 |
| | | 28 |
| | Results..... | 29 |
| | | |
| | PRISMA Flow chart | 29 |
| | Forest Plot depicting overall BPCR status in India | 31 |
| | Forest Plot depicting (weight of studies) | 33 |
| | Forest Plot depicting subgroup analysis-Delivery status | 35 |
| | wise | |
| | Forest Plot depicting subgroup analysis-Region wise | 35 |
| | Prevalence of components of BP and CR | 38 |
| | Funnel Plot | 40 |
| | Funnel Plot after performing Eggers test | 40 |
| | Graph depicting;Quality Assessment | 41 |
| | Discussion..... | 43 |
| | | |
| | Limitation..... | 47 |
| | | |
| | Conclusion | 49 |
| | | |
| | Reference..... | 51 |
| | | |

Acronyms/Abbreviation-

BPCR- Birth Preparedness and Complication Readiness

GoI- Government of India

JBI- Joanna Briggs Institute

JHPIEGO- Johns Hopkins Program for International Education in Gynaecology and Obstetrics

JSY- Janani Suraksha Yojana

JSSK- Janani Shishu Suraksha Karyakaram

LaQshya- Labour room Quality Improvement Initiative

MMR- Maternal Mortality Ratio

MMR- Maternal Mortality Rate

PMSMA- Pradhan Mantri Surakshit Matritva Abhiyan

PRISMA- Preferred Reporting Items for Systematic Review and Meta-Analysis

SRMA- Systematic review and meta-analysis

WHO-World Health Organization

ABOUT IIHMR DELHI

The International Institute of Health Management Research (IIHMR), New Delhi is allied to the ‘Society for Indian Institute of Health Management Research’ which was established in October 1984 under the Societies Registration Act-1958. IIHMR-Delhi was setup in 2008 in response to the growing needs of sustainable management and administration solutions critical to the optimal function of healthcare sector both in India and in the Asia-Pacific region.

We are a leading institute of higher learning that promotes and conducts research in health and hospital management; lends technical expertise to policy analysis and formulation; develops effective strategies and facilitates efficient implementation; enhances human and institutional capacity to build a competent and responsive healthcare sector. Our multi-dimensional approach to capacity building is not limited to academic programs but offers management development programs, knowledge and skills-based training courses, seminars/webinars, workshops, and research studies. Our four core activities are...

- Academic courses at masters and doctoral level in health and hospital management to meet the growing need of skilled healthcare professionals.
- Research that has high relevance to health policies and programs at national and global level.
- Continued education through management development programs and executive programs for working professionals to help them upgrade their knowledge and skills in response to the emerging needs of the industry.
- Technical consultation to the national and state-level flagship programs to address the gaps in planning as well as implementation.

International Institute of Health Management Research, New Delhi (IIHMR-Delhi)

Over the years IIHMR-Delhi has emerged as an institute of repute both nationally and globally for producing socially conscious, skilled and vibrant top-class health care management professionals. Our graduates are well-matched for the ever-changing health care sector and evolving social milieu. The institute has progressed as a leader in research, teaching, training, community extension programmes and policy advocacy in the field of health care. IIHMR has carved out a niche for itself through its cutting-edge academic curriculum, infrastructure, accomplished multi-disciplinary faculty and research.

The Institute as an autonomous body of international stature has been developing leaders for several years to shape tomorrow's healthcare by equipping the students in the fields of health, hospital, and health information technology. The Institute's dynamic health care research programmes provide rigorous training in management, health systems, hospital administration, health care financing, economics, and information technology

Commitment to Inclusive Excellence

As an institute, IIHMR-Delhi is committed to creating an environment of higher learning that can serve as the model for the kind of society it strives to build – one of equity, social justice and mutual support. We have also made a concerted effort to promote the ethos and philosophies amongst today's students and nurture them into growing as effective managers, to think both critically and ethically, to learn to cope with ethical dilemmas and apply systems-thinking approaches to serious and complex societal problems. Our internationally renowned faculty lead multidisciplinary health research in multifarious areas such as public health, health services, health economics, hospital management, social determinants of health, mental Health and other topics of global and national interest.

The IIHMR is invited by various governmental and civil society organizations to provide technical support for capacity building and policy research needs that culminates in developing innovative and equitable health care strategies and provide advocacy support for health policy and planning. The institute also responds to the global health threats, natural disasters, conflict and related humanitarian crisis. In addition to the Masters and doctoral level programmes, IIHMR-D also offers several highly specialized and popular Management Development Programmes (MDP) to wide range of health professional in the country and overseas which largely addresses educational needs amongst in-service aspirants.

INTRODUCTION-

The health of pregnant women, from conception to the postnatal period, must be treated as a priority. Each stage of the process should be a positive experience, enabling both mother and baby to realize their full potential for health and happiness.

In 2012, a total of 213 million pregnancies took place, with the majority (89%) occurring in developing nations and a smaller proportion (11%) in the developed world. It highlights the disparities in access to quality health services.(1) Whereas, In India from 2015 to 2019, an estimated 48.5 million pregnancies occurred each year.(2)

Maternal Mortality can be defined as the number of female deaths per annum due to any cause related to or caused by pregnancy, childbirth, or up to 42 days after the termination of the pregnancy, regardless of the length and location of the pregnancy, is recorded.(3)

According to the WHO and a Study by (Say et al. 2014) Suggests, that approximately three-quarters (75%) of maternal fatalities are due to direct obstetric causes that can be attributed to severe haemorrhage, infection typically occurring following childbirth, high blood pressure during gestation (pre-eclampsia and eclampsia), embolism, unsafe abortion and rest (25%) deaths was due to indirect obstetric causes of pregnancy resulting from previously existing disease like cardiovascular diseases, HIV, severe anaemia, diabetes and hepatitis when aggravated by Pregnancy.(4–7)

Although some advances have been made in the past two decades, the number of women who died in 2020 due to pregnancy and childbirth-related causes is still exceedingly high at 287,000, which is not an acceptable figure.(8)

In 2020, virtually 95% of maternal fatalities took place in lower and lower-middle-income countries (LMIC), with most of them being avoidable. Sub-Saharan Africa and Southern

Asia were the locations of 87% (253,000) of the estimated global maternal deaths. Between 2000 and 2020, Eastern Europe and Southern Asia experienced the most drastic decrease in the maternal mortality ratio (MMR): a reduction of 70% (from 38 MMR to 11) and 67% (from 408 MMR to 134) respectively.(4)

The Indian Government has made significant progress in reducing MMR from 103 to 97 per lakh live births between 2017-2019 and 2018-2020 respectively, surpassing the goal set by the NHP 2017 for 2020. However, the MMR still needs to be improved further to reach the Sustainable Development Goal of 70 per 100,000 live births by 2030.(9,10)

To reduce the maternal mortality rate (MMR) in India, strategies such as prioritizing maternal and child health (MCH) services, ensuring safe delivery and postpartum periods, providing quality primary health care, promoting health education and reproductive behaviour and rights of safe motherhood should be adopted (11) and as well as the proper management of potential delays.(12,13)

To ensure maternal health, it is essential that all women, including adolescents, have access to contraception, safe abortion services, and quality post-abortion care. Furthermore, pregnant women should receive quality care during and after childbirth and should be attended by skilled health professionals. Additionally, measures such as the timely administration of oxytocic, good hygiene, early detection and treatment of infections, and the use of drugs such as magnesium sulphate for pre-eclampsia should be adopted to reduce the risk of maternal mortality.(4)

Thaddeus and Maine (14) have identified three delays associated with receiving care during childbirth: the decision to seek care, identifying and reaching a medical facility, and receiving adequate and appropriate treatment. To address these delays, Johns Hopkins Program for International Education in Gynaecology and Obstetrics (JHPIEGO) has created the Birth Preparedness and Complication Readiness (BPCR) concept to

ensure that expectant mothers are able to access the care they need without unnecessary delays.(15,16)




Figure 1:- Conceptual Diagram of How BPCR may increase the use of skilled care

BPCR approach is an effective way to ensure the use of skilled maternal and neonatal services in a timely manner.(17) Having a BPCR plan in place helps to ensure the best possible outcome by being prepared for any unexpected issues that may arise during the birthing process.(18) A study conducted in Ethiopia revealed that the Birth Preparedness and Complication Readiness (BPCR) had a considerable impact on the utilization of

skilled care. Furthermore, a meta-analysis of 14 randomized trials demonstrated that, given adequate coverage of the population, BPCR interventions are effective in lowering the maternal and mortality rate in areas with limited resources.(19–21)

BPCR helps to speed up the decision to seek care in two ways. Firstly, BP encourages people to plan to have a trained professional at the time of childbirth. If this plan is put into action, the woman can get the necessary care before any potential issues arise during the birth, thus avoiding the two delays. Secondly, CR raises the awareness of danger signs among families and communities, improving the recognition of the problem and speeding up the decision to seek care.(15,22,23)

The Government of India (GoI) launched several flagship programmes under RMNCH+A some of them are - Janani Suraksha Yojana (JSY), a demand promotion and conditional cash transfer scheme which strives to reduce Maternal and Infant Mortality by encouraging institutional delivery among poor pregnant women. Janani Shishu Suraksha Karyakaram (JSSK) to provide free delivery services, drugs, diagnostics, blood, diet, transport, and treatment for newborns in public healthcare institutions. Pradhan Mantri Surakshit Matritva Abhiyan (PMSMA) launched to ensure quality antenatal care and high-risk pregnancy detection on the 9th of every month. LaQshya, a Labour room Quality improvement Initiative, recently introduced to further accelerate the decline in MMR by strengthening key processes related to labour rooms and maternity operation theatres, as well as ensuring Respectful Maternity Care.(24) There is no specific Program in India which focuses specifically on BPCR, but the PMSMA, consists of one objective, related to BPCR, it states – “appropriate BPCR provided for each pregnant woman, especially those who are identified as having a risk factor or co-morbid condition.”(25)

Rationale-

To date, no systematic review and meta-analysis have been conducted on BPCR in India. This systematic review and meta-analysis (SRMA) will provide a comprehensive overview of the current situation of BPCR in India. This analysis will bring to light the gaps in the existing knowledge base, identify potential areas of improvement, and inform evidence-based strategies to improve BPCR in India.

METHODOLOGY-

Study Registration-

This Systematic Review adhered to the Preferred Reporting Items for Systematic Review and Meta-Analysis (PRISMA) guidelines (26) and the protocol was registered at the International Prospective Register of Systematic Reviews (27) (registration ID: CRD42023396109). (Annexure 1)

Search strategy –

Keywords were used to create a search strategy for addressing the research questions. Systematic search was performed by combining every feasible sequence of all the categories of keywords. The Medical Subject Headings (MeSH) terms and truncated keywords were mixed using the relevant Boolean logic operators i.e., AND, OR, and NOT. The authors (BT, TS, AP, DG) pretested the search strategy to ensure appropriateness of the search strategy in retrieving the relevant articles and subsequent modifications.

Based on the inclusion criteria, review was done on the following search engines and other sources. Major databases like; PubMed, Google Scholar, Cochrane Library, and Proquest were used to review published studies. The reference lists of already identified studies were also searched to retrieve additional articles. All published articles up to January 31st, 2023, were included in this review. The search was done using the following search terms using PICO format given in table 1.

Table 1: PICO Format

| | |
|---------------------|---|
| <i>Population</i> | Pregnant women: "pregnan*"[Title/Abstract] OR "pregnant women"[Title/Abstract] OR "antenatal"[Title/Abstract] OR "anc"[Title/Abstract] |
| <i>Intervention</i> | - |
| <i>Comparison</i> | - |
| <i>Outcome</i> | Birth preparedness: "birth preparedness"[Title/Abstract] OR "preparedness"[Title/Abstract] OR "preparing for birth"[Title/Abstract] OR "emergency preparedness"[Title/Abstract] OR "birth plan"[Title/Abstract] |
| | Complication Readiness: "danger signs"[Title/Abstract] OR "readiness"[Title/Abstract] OR (("recognisable"[All Fields] OR "recognise"[All Fields] OR "recognised"[All Fields] OR "recognises"[All Fields] OR "recognising"[All Fields] OR "recognize"[All Fields] OR "recognized"[All Fields] OR "recognizes"[All Fields] OR "recognizing"[All Fields]) AND "danger sign"[Title/Abstract]) OR "obstetric complication"[Title/Abstract] OR "pregnancy complication"[Title/Abstract] OR "obstetric danger sign"[Title/Abstract] OR "maternal complications"[Title/Abstract] OR "maternal health"[Title/Abstract] OR "newborn health"[Title/Abstract] |

*I and C are not mentioned as only descriptive studies were included in the SRMA.

Study selection and eligibility criteria-

Inclusion criteria

This review included cross-sectional studies conducted at the community or institutional level on BPCR in India among pregnant and recently delivered women in India with BPCR as outcome. All studies in English language irrespective of the time of data collection or publication year were included.

Exclusion criteria

Studies which did not reported the outcome variable as BPCR, didn't specify study population, qualitative studies were excluded.

Selection process-

Studies which met the stated inclusion criteria were retrieved and reviewed by two reviewers independently (TS and BT). The two reviewers independently evaluated the eligibility of the studies obtained from the literature searches for title and abstract screening.

“Rayyan.ai” software, a web-based automated screening tool which uses text mining methods to facilitate semi-automatic screening of records for SRs for title and abstract screening. Records were screened and labelled for either inclusion, exclusion, or “maybe” relevant to the subject of the review, and also highlighted the words for inclusion and exclusion (which significantly assisted manual screening), and both the reviewers choose the reason(s) for excluding a record by turning ON the blinding mode.

The generated reports from the two reviewers were retrieved and selected articles will then undergo a full-text screening by (TS and BT). In cases of discrepancies, an agreement was reached by consensus with the advice of third-member arbitrators (AKP and DG).

Justification was given for excluding the studies with proper reason. The search process was presented in the form of a PRISMA flow chart.

Data extraction and data collection-

Two reviewers (TS and BT) independently extracted data using a predefined data extraction Microsoft excel spreadsheet. The sheet included questions on the author's name, publication year, study design, sample size, study area, participants, mean age of participants, response rate and prevalence of BPCR. Additionally, the tool contains information on; percentage of women who saved money for birth and emergency case, women who prepared blood donor, women who identified skilled birth attendant, women who were aware of danger signs during pregnancy, labour, postpartum, and newborn, women who arranged transportation, women who identified place of birth, women who planned health facility delivery, women who knew about financial assistance provided by government in JSY, and women who knew about transportation provided by government in JSSK.

Main Outcome-

The main outcome of this review was to know the BPCR among pregnant and recently delivered women in India.

Measures of effect-

Prevalence of BPCR among pregnant and recently delivered women in India.

Risk of Bias (ROB) Quality Assessment-

All the studies will be assessed for methodological quality, risk of bias to check the validity of the study findings before identifying and finalizing the study for use. Two reviewers (TS and BT) assessed the risk of bias of the included studies independently by using the Joanna Briggs Institute (JBI)(26) critical appraisal checklist, which is devised for studies reporting prevalence data. The JBI methodology checklist consists of 9 questions and is categorized into “Yes”, “NO”, “Unclear” and “Not applicable”.

Overall risk assessment was done by giving each study with a score of 0-9 and categorising them into any of the three categories: - (0-3=HIGH, 4-6=MEDIUM, and 7-9=LOW). After final assessment it was found that Most of the studies have low risk of Bias. Any disagreements over the assessment of the risk of bias and research quality between two reviewers was settled by involving the third reviewer author.

Data synthesis, statistical analysis, and investigation of heterogeneity-

The data was analysed using **CMA (Comprehensive meta-analysis tool) and R studio software**. The analysis was done by **AP, DG, TS, and BT**. We performed a meta-analysis applying fixed and random effects model, considering the assessed heterogeneity between the studies. The pooled estimates were reported by all the studies and the findings were presented using forest plots. Statistical heterogeneity was assessed using a combination of visual inspection of the forest plot along with consideration of the Cochran's Q test, and the I^2 statistic results, in reference to the Cochrane Handbook Criteria. A probability value of $p < 0.05$ was taken to indicate statistically significant heterogeneity. When the results fell below 25%, between 25% and 75%, and over 75%, respectively, heterogeneity was deemed low, moderate, and high respectively. When discrepancies between study outcomes go beyond those only due to chance, statistical heterogeneity was present. Funnel plot and Egger's test were applied to explore the

possibility of the publication bias in the studies. Sensitivity analyses was conducted to assess the influence of individual studies on the pooled estimates and to evaluate whether the overall estimates were dominated by one single study.

Subgroup analysis-

As per the data availability, we have done a subgroup analysis for BPCR by delivery status (recently delivered and pregnant women) and geographical region wise (East, West, North, South, and Central). (27)

Definition of BPCR-

BPCR is a strategy to promote the timely use of skilled maternal and neonatal care, especially during childbirth, based on the theory that preparing for childbirth and being ready for complications reduces delays in obtaining this care.(28) BPCR is measured by key elements. The key elements included arrangement for transportation, saving money for delivery, identify skilled birth attendant, identifying place of delivery and identifying blood donor for the case of emergency, knowledge regarding danger signs during pregnancy, labour, postpartum and newborn. All studies that used the above definition of BPCR were included in this review.(15)

RESULTS

Authors have identified records from major databases like PubMed, Cochrane, Proquest (n=1349) records in total and from other sources identified (n=42) records by reference/citation search. After removing (n=378) records before screening, (n=971) articles were finalised to undergo screening, where (n=958) records were removed because of duplication and (n=42) articles from other sources were screened for title and abstract (**Figure 2**).

After title and abstract screening (n=13) articles from major databases and (n=23) from other searches were eligible for full text screening. Lastly (n=36) articles were eligible to be included, which undergone full-text screening and out of which (n=1) article was excluded because of no full text and Finally (n=35) Full text articles were included.

DESCRIPTION OF THE STUDIES

PRISMA Flow diagram for Systematic Reviews, 2020




Figure 2:- PRISMA Flow chart summarizing the article selection process

The forest plot (**figure 3**) showed results of meta-analysis of 27 studies that assessed the proportion of pregnant women who are aware of BPCR in different Regions of India. The overall pooled proportion is 0.49 (95% CI: 0.43 to 0.55), which means that on average, about half of the pregnant women are aware of BPCR across. However, there is a high and significant degree of heterogeneity among the studies ($I^2 = 94\%$, $\tau^2 = 0.3110$, $p < 0.01$), which meant that the studies are not consistent and may have different sources of variation. The study with the lowest proportions is Vishwanathan VT (2020) with 0.15 (95% CI: 0.12 to 0.19), while the study with the highest proportion is Akshaya KM with 0.79 (95% CI: 0.72 to 0.84). The CIs of most study overlap with each other and with the overall estimate, which means that there is no significant difference between them in terms of BPCR awareness. However, some study had CIs that do not include the pooled proportion, such as Gurung J (2017), Indira NC (2021), Kushwah SS (2009), and Patil MS (2016), which meant that they were significantly different from the overall estimate and may have some special characteristics or factors that influence their BPCR awareness. The diamond for the random effects model is also slightly to the right of the line of no effect, indicating a positive and statistically significant pooled proportion of BPCR across all studies. The point estimate is 0.50, with a 95% confidence interval of [0.37, 0.63]. This meant that if each study had its own true proportion of BPCR that varied around an overall mean, we would expect it to be between 37% and 63% with 95% probability.




Figure 3:- Forest Plot Depicting Overall BPCR status in India

Forest Plot (**Figure 4**) depicted weight of each study which was analyzed through random and common effect model. We considered Random effect Model for assessing the studies as it assumed that there were different true effects for each study and the difference among the study is due to chance and real variation therefore it gives balanced weight to all studies.




Figure-4 Forest plot Depicting (Weight of Studies).

In a study by Kushwah SS (2009) has a total sample size of 2022 so, its weight in Common Effect model was 20.9% which is very high from other studies, whereas in random effect model the same study has a weight of 4.1% which shows the equal balance of weights among all studies.

Subgroup analysis taking delivery status into account was performed. The overall pooled proportion of knowledge of BPCR in RDW was 0.54 (95% CI: 0.46 to 0.62) and in PW was 0.50 (95% CI:0.45 to 0.55) with high heterogeneity ($I^2= 91%$, Tau Square= 0.1221 and with a p- value < 0.01 which is statistically significant. As the CIs among the subgroups overlap there was no statistical significance observed. Same happened when the subgroup analysis was done region wise. The overall pooled proportion of knowledge of BPCR in East region is 0.47 (95% CI: 0.38 to 0.56), North region is 0.47 (95% CI: 0.36 to 0.57), Central region is 0.46 (95% CI: 0.43 to 0.49), West region is 0.48 (95% CI: 0.26 to 0.70), South region is 0.56 (95% CI: 0.45 to 0.67). **(Figure 5 and 6)**




Figure 5: - Forest plot Depicting subgroup analysis – Delivery Status wise




Figure 6:- Forest plot Depicting Sub group analysis – Region wise

Prevalence of Components of BP and CR

From the 35 Studies that were finalized, all of them were divided their BPCR indicators into two main heads i.e, Knowledge regarding BP and Knowledge regarding CR (Table 1).

Several Components of danger signs during pregnancy among them 13 studies have reported at least one danger sign of pregnancy and the lowest prevalence was found (9.3%) to be in studies reporting three or more danger signs of postpartum. Knowledge regarding danger signs during pregnancy were reported by 10 studies with a prevalence of 34.9%, $i^2=97.4$ and p value=0.00 and similarly, prevalence among other components like Knowledge regarding danger signs during labor, postpartum, and in neonate was found to be 11.6%, 12.2% and 18.8% and the i^2 was 98%, 91.1% and 95.3%, which depicts high heterogeneity.

Similarly, components of BP were also assessed by different studies among which 14 studies reports that the prevalence of women who identified a place of delivery was 59.4% and 13 studies report that prevalence of women who got financial assistance by JSY scheme was 55.5%.

BP component which was found similar in most of the studies was women who have identified transport and saved money for their delivery and the prevalence of both the components was found to be 46% and the i^2 was 98.8, with a p -value of 0.00.

The highest prevalence 86.7% was seen in women who attended their 1st antenatal visit with a skilled person during 1st trimester and the least prevalence 10.1% was seen in women who identified blood donor.

| | Components | Events | Sample size | Prevalence | I2 | TauSq | P value | number of studies | LL | UL |
|--|--|--------|-------------|--------------|------|-------|---------|-------------------|------|------|
| Knowledge regarding Danger signs | Knowledge regarding danger signs during pregnancy | 754 | 2325 | 34.9% | 97.4 | 0.95 | 0.00 | 10 | 0.22 | 0.50 |
| | Knowledge regarding danger signs during labor | 237 | 1553 | 11.6% | 98.0 | 1.69 | 0.00 | 5 | 0.04 | 0.29 |
| | Knowledge regarding danger signs during postpartum | 267 | 1687 | 12.2% | 91.1 | 0.32 | 0.00 | 6 | 0.08 | 0.19 |
| | Knowledge regarding danger signs in neonate | 195 | 1189 | 18.8% | 95.3 | 0.57 | 0.00 | 4 | 0.10 | 0.33 |
| | 3 or more ds of newborn care | 430 | 1512 | 12.3% | 97.6 | 0.88 | 0.00 | 4 | 0.05 | 0.27 |
| | 3 or more ds of Postpartum | 169 | 1512 | 9.3% | 94.2 | 0.66 | 0.00 | 4 | 0.04 | 0.20 |
| | 3 or more ds of labor n childbirth | 322 | 1512 | 15.7% | 92.7 | 0.30 | 0.00 | 4 | 0.09 | 0.26 |
| | 3 or more ds of preg | 423 | 1512 | 22.0% | 93.6 | 0.26 | 0.00 | 4 | 0.14 | 0.33 |
| | 1 ds of newborn | 837 | 2770 | 28.8% | 98.5 | 1.31 | 0.00 | 8 | 0.15 | 0.47 |
| | 1 ds of postpartum | 567 | 1863 | 30.4% | 97.6 | 1.41 | 0.00 | 6 | 0.14 | 0.54 |
| | 1 ds of labor | 649 | 2367 | 30.0% | 97.7 | 1.04 | 0.00 | 6 | 0.16 | 0.50 |
| | 1 DS of Pregnancy | 1559 | 4339 | 38.7% | 98.7 | 1.59 | 0.00 | 13 | 0.24 | 0.56 |
| Knowledge regarding Birth preparedness | place of delivery | 2192 | 5964 | 59.4% | 99.1 | 2.50 | 0.00 | 14 | 0.39 | 0.77 |
| | transport jssk jsy | 1895 | 5286 | 50.6% | 99.0 | 1.32 | 0.00 | 11 | 0.34 | 0.67 |
| | Financial assistance jsy | 3276 | 5566 | 55.5% | 98.7 | 1.06 | 0.00 | 13 | 0.41 | 0.69 |
| | skilled birth attendant for deli | 6183 | 8345 | 83.3% | 98.6 | 1.86 | 0.00 | 22 | 0.73 | 0.90 |
| | identify blood donor | 1008 | 7842 | 10.1% | 98.0 | 1.45 | 0.00 | 19 | 0.06 | 0.16 |
| | identify transport | 6396 | 14232 | 46.0% | 98.8 | 1.28 | 0.00 | 34 | 0.37 | 0.56 |
| | Saved Money | 5839 | 13774 | 46.0% | 98.8 | 1.28 | 0.00 | 33 | 0.40 | 0.56 |
| | Four or more ANC's | 2205 | 4910 | 51.5% | 98.8 | 1.27 | 0.00 | 13 | 0.36 | 0.66 |
| | 1st antenatal visit with a skilled person during 1st trimester | 1958 | 2283 | 86.7% | 96.1 | 0.71 | 0.00 | 7 | 0.77 | 0.93 |
| registration within 12 weeks | 2028 | 3511 | 63.6% | 99.0 | 1.37 | 0.00 | 8 | 0.44 | 0.80 | |

Table 2:-Prevalence of components of BP and CR

PUBLICATION BIAS:

To detect publication bias we constructed funnel plot in which most of the studies were lying in the middle of the plot which showed that there is no major variance among the studies. **(Figure 8)** To statistically prove that there was no major variance we used Egger's Test, which states that if the p-value < 0.05 , it indicates evidence of publication bias, and in our study, we found out that the p value is 0.59 ($p > 0.05$). **(Figure 9)**




Figure 7:- Funnel Plot




Figure 8:- Funnel Plot after Performing Eggers Test.

Graph depicting: Quality Assessment

The author's estimated ROB which was found to be low for 28 studies, unclear for 7 studies, and high for 0 studies. Detailed information regarding the results of the quality assessment of the selected studies is provided in **Figure 10**.




Figure 10:- Stacked bar chart depicting ROB.

DISCUSSION

The overall BPCR score of 49% in our study indicates that there is still a gap in the utilization of BPCR services among pregnant women in India. This score is slightly higher than the 44.9% reported by Girma et al. (2013)(29) in a similar study in Ethiopia. The variation in the BPCR scores may be attributed to the differences in the socio-demographic characteristics, health system factors, and cultural practices of the study populations. Compared to other studies in Nigeria, Ethiopia, and India by Akinwaare et al. (2015)(30), Gedefa et al. (2017)(31), and Nimavat et al. (2018)(32), our study had a moderate BPCR score of 49%, which ranged from 30.6% to 58.7% in these studies. Therefore, there is a need to improve the awareness and utilization of BPCR services among pregnant women in our setting, as well as to address the barriers and facilitators that influence their decision-making and behavior regarding BPCR.

The prevalence of women who saved money for their delivery was found to be 46% in our study, which is an important indicator of BPCR. This prevalence is higher than the 38.7% reported by Berhe et al. (2016)(33) in a study in Ethiopia, but lower than the 63.4% reported by Akinwaare et al. (2015)(30) in a study in Nigeria, the 59.6% reported by Nimavat et al. (2018)(32) in a study in India, and the 83.3% reported by Moran et al. (2018)(34) in a study in Burkina Faso. The differences in the prevalence of saving money for delivery may reflect the variations in the economic status, access to financial services, and social norms of the study populations. Saving money for delivery can help women to overcome the financial barriers to access skilled care during pregnancy and childbirth, and to cope with any unforeseen complications that may arise. Therefore, there is a need to promote and facilitate saving money for delivery among pregnant women in our setting, as well as to provide them with adequate information and counselling on the benefits and options of saving money for delivery.

The prevalence of women who have knowledge of key danger signs of pregnancy was 34.9% in our study, which is a low level of awareness that can compromise the timely recognition and management of pregnancy complications. This prevalence is higher than the 26.3% reported by Berhe et al. (2016)(33) in a study in Ethiopia, but lower than the 52% reported by Akinwaare et al(30). (2015) in a study in Nigeria, the 83.3% reported by Moran et al. (2018)(34) in a study in burkina faso, and the 42% reported by Mukhopadhyay et al. (2016)(28) in a study in India. The differences in the prevalence of knowledge of key danger signs of pregnancy may be related to the variations in the educational level, exposure to mass media, and quality of antenatal care services of the study populations. Knowledge of key danger signs of pregnancy is essential for pregnant women to seek prompt and appropriate care when they experience any signs of potential complications, and to prevent maternal and neonatal morbidity and mortality. Therefore, there is a need to improve the knowledge of key danger signs of pregnancy among pregnant women in our setting, as well as to provide them with effective health education and counseling on the identification and management of key danger signs of pregnancy.

The prevalence of women who have arranged transport for their delivery was 46% in our study, which is a moderate level of preparedness that can facilitate the access to skilled care during childbirth. This prevalence is similar to the 46.1% reported by Moran et al. (2018)(34) in a study in Burkina Faso, but varies from the other studies in Ethiopia, Nigeria, and India by Berhe et al. (2016), Akinwaare et al. (2015), Nimavat et al. (2018), and Mukhopadhyay et al. (2016),(28,30,32,33) which reported prevalences ranging from 20.59% to 58.6%. The differences in the prevalence of arranging transport for delivery may depend on the availability, affordability, and acceptability of transport services in the study settings. Arranging transport for delivery can help women to overcome the geographical and financial barriers to reach a health facility in time, and to avoid delays and complications during delivery. Therefore, there is a need to encourage and support

pregnant women in our setting to arrange transport for their delivery, as well as to improve the transport system and infrastructure in our area.

The prevalence of women who have identified a place of birth for their delivery was 59.4% in our study, which is a relatively high level of preparedness that can influence the choice and utilization of skilled care during childbirth. This prevalence is slightly higher than the 54.85% reported by Berhe et al. (2016)(33) in a study in Ethiopia, which had a similar study design and population. The similarity in the prevalence of identifying a place of birth for delivery may reflect the common socio-cultural and health system factors that affect the decision-making and behavior of pregnant women in both settings. Identifying a place of birth for delivery can help women to plan ahead and to select a health facility that meets their needs and preferences, and to avoid uncertainty and confusion during delivery. Therefore, there is a need to maintain and enhance the prevalence of identifying a place of birth for delivery among pregnant women in our setting, as well as to provide them with adequate information and counseling on the benefits and options of identifying a place of birth for delivery.

The prevalence of women who have arranged a blood donor for their delivery was 10.1% in our study, which is a very low level of preparedness that can increase the risk of maternal and neonatal mortality due to hemorrhage. This prevalence is similar to the 8.18% reported by Berhe et al. (2016)(33) in a study in Ethiopia, and the 9.9% reported by Mukhopadhyay et al. (2016)(28) in a study in India, but higher than the 2.7% reported by Nimavat et al. (2018)(32) in another study in India. The similarity in the prevalence of arranging a blood donor for delivery may indicate the common challenges and barriers that pregnant women face in accessing and obtaining blood transfusion services in these settings. Arranging a blood donor for delivery can help women to cope with any potential complications that may require blood transfusion, and to save lives during delivery. Therefore, there is a need to improve the prevalence of arranging a blood donor for

delivery among pregnant women in our setting, as well as to improve the availability, accessibility, and quality of blood transfusion services in our area.

LIMITATION-

The Meta- analysis did not address the factors affecting for the low Utilization of BPCR, therefore Researchers should Conduct further studies in this Area.

CONCLUSION-

Our study highlights the low prevalence of BPCR in India and the factors associated with it. Our findings underscore the need for targeted interventions to improve BPCR and suggest that there is a need to improve the awareness and utilization of BPCR services among pregnant women in India, as well as to address the factors that influence their decision-making and behavior regarding BPCR. India's health system is already overburdened by adverse health outcomes. Therefore, scaling up cost-effective interventions like BPCR that have a positive overall effect is necessary. Our study suggests that birth preparedness and complication readiness (BPCR) is such an intervention that should be given utmost importance to reduce maternal morbidity and mortality and to achieve the Sustainable Development Goals.

References:-

- 1.Sedgh G, Singh S, Hussain R. Intended and Unintended Pregnancies Worldwide in 2012 and Recent Trends. *Stud Fam Plann* [Internet]. 2014 Sep [cited 2023 Apr 25];45(3):301–14. Available from: <https://onlinelibrary.wiley.com/doi/10.1111/j.1728-4465.2014.00393.x>
- 2.India [Internet]. Guttmacher Institute. 2022 [cited 2023 Apr 25]. Available from: <https://www.guttmacher.org/regions/asia/india>
- 3.Maternal deaths [Internet]. World Health Organization; 2023. Available from: <https://www.who.int/data/gho/indicator-metadata-registry/imr-details/4622>
- 4.Maternal mortality [Internet]. [cited 2023 Apr 5]. Available from: <https://www.who.int/news-room/fact-sheets/detail/maternal-mortality>
- 5.Say L, Chou D, Gemmill A, Tunçalp Ö, Moller AB, Daniels J, et al. Global causes of maternal death: a WHO systematic analysis. *Lancet Glob Health* [Internet]. 2014 Jun [cited 2023 Apr 25];2(6):e323–33. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S2214109X1470227X>
- 6.Gupta S. Study of Direct and Indirect Causes of Maternal Mortality: A Study from Tertiary Care Centre of Bhopal. *Sch Int J Obstet Gynecol*.
- 7.Storm F, Agampodi S, Eddleston M, Sørensen JB, Konradsen F, Rheinländer T. Indirect causes of maternal death. *Lancet Glob Health* [Internet]. 2014 Oct 1 [cited 2023 Apr 25];2(10):e566. Available from: [https://www.thelancet.com/journals/langlo/article/PIIS2214-109X\(14\)70297-9/fulltext](https://www.thelancet.com/journals/langlo/article/PIIS2214-109X(14)70297-9/fulltext)

8. Maternal health [Internet]. [cited 2023 Apr 8]. Available from: <https://www.who.int/health-topics/maternal-health>
9. Ministry of Women and Child Development. Significant Decline in Maternal Mortality in India [Internet]. Ministry of Women and Child Development; 2022. Available from: [https://pib.gov.in/FeaturesDeatils.aspx?NoteId=151238&ModuleId%20=%202#:~:text=MMR%20in%20the%20country%20declined,MMR%20of%20211%20\(2017\).](https://pib.gov.in/FeaturesDeatils.aspx?NoteId=151238&ModuleId%20=%202#:~:text=MMR%20in%20the%20country%20declined,MMR%20of%20211%20(2017).)
10. SRS - Maternal Mortality Bulletin | Government of India [Internet]. [cited 2023 Apr 5]. Available from: <https://censusindia.gov.in/census.website/data/SRSMMB>
11. Prakash A, Swain S, Seth A. Maternal mortality in India: current status and strategies for reduction. *Indian Pediatr.* 1991 Dec;28(12):1395–400.
12. Blencowe H, Cousens S, Jassir FB, Say L, Chou D, Mathers C, et al. National, regional, and worldwide estimates of stillbirth rates in 2015, with trends from 2000: a systematic analysis. *Lancet Glob Health.* 2016 Feb;4(2):e98–108.
13. Kar M, Karmee N, Satapathy DM. Birth preparedness and complication readiness among pregnant and recently delivered women in villages of a block of Ganjam, Odisha, India: a community based cross-sectional study. *Int J Reprod Contracept Obstet Gynecol* [Internet]. 2019 Apr 29 [cited 2023 Apr 6];8(5):2003. Available from: <https://www.ijrcog.org/index.php/ijrcog/article/view/6417>
14. Thaddeus S, Maine D. Too far to walk: Maternal mortality in context. *Soc Sci Med* [Internet]. 1994 Apr [cited 2023 Apr 6];38(8):1091–110. Available from: <https://linkinghub.elsevier.com/retrieve/pii/0277953694902267>

15. Barco RCD. JHPIEGO Brown's Wharf 1615 Thames Street Baltimore, Maryland 21231-3492, USA. Fam CARE [Internet]. Available from: https://pdf.usaid.gov/pdf_docs/pnada619.pdf
16. Salroo FN, Nazir ST, Gadoo MM. BIRTH PREPAREDNESS AND COMPLICATION READINESS AMONG PREGNANT WOMEN ATTENDING A MATERNAL AND CHILD CARE HOSPITAL OF GOVERNMENT MEDICAL COLLEGE IN SOUTH KASHMIR, INDIA: A CROSS-SECTIONAL STUDY. Asian J Pharm Clin Res [Internet]. 2022 Jan 7 [cited 2023 Apr 6];35–40. Available from: <https://innovareacademics.in/journals/index.php/ajpcr/article/view/46414>
17. Berhe AK, Muche AA, Fekadu GA, Kassa GM. Birth preparedness and complication readiness among pregnant women in Ethiopia: a systematic review and Meta-analysis. Reprod Health [Internet]. 2018 Dec [cited 2023 Apr 8];15(1):182. Available from: <https://reproductive-health-journal.biomedcentral.com/articles/10.1186/s12978-018-0624-2>
18. Pandey P, Srivastava R, Kumari K, Pandey M. Status of Birth Preparedness and Complication Readiness of Pregnant Women and Recently Delivered Women in Rural Varanasi: Assessment of Current Scenario. Indian J Community Med [Internet]. 2022 [cited 2023 Apr 6];47(2):249. Available from: <http://www.ijcm.org.in/text.asp?2022/47/2/249/350342>
19. Soubeiga D, Gauvin L, Hatem MA, Johri M. Birth Preparedness and Complication Readiness (BPCR) interventions to reduce maternal and neonatal mortality in developing countries: systematic review and meta-analysis. BMC Pregnancy Childbirth [Internet]. 2014 Apr 4 [cited 2023 Apr 9];14:129. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4234142/>

20. Agarwal S, Sethi V, Srivastava K, Jha PK, Baqui AH. Birth Preparedness and Complication Readiness among Slum Women in Indore City, India. *J Health Popul Nutr* [Internet]. 2010 Sep 6 [cited 2023 Apr 6];28(4):383–91. Available from: <http://banglajol.info/index.php/JHPN/article/view/6045>
21. Karir DS, Panda DB, Hota DG, Kumari DS, Panda DPC. Birth Preparedness and Complication Readiness among Recently Delivered Mothers-A Cross Sectional Survey in an Urban Community in Eastern India. *Clin Med*. 2022;09(02).
22. Patel NA, Mehta JP, Unadkat SV, Yadav SB. Birth preparedness: studying its effectiveness in improving maternal health in urban slums of Jamnagar, Gujarat. *Int J Community Med Public Health* [Internet]. 2017 Nov 23 [cited 2023 Apr 6];4(12):4569. Available from: <http://www.ijcmph.com/index.php/ijcmph/article/view/2096>
23. Vidhyashree MD, S. J, P. K, V. L, S. S, K. S. Birth preparedness and complication readiness for a safe motherhood among antenatal women attending an urban health centre, Pudupet. *Int J Community Med Public Health* [Internet]. 2020 Oct 26 [cited 2023 Apr 6];7(11):4345. Available from: <https://www.ijcmph.com/index.php/ijcmph/article/view/6931>
24. Maternal Health :: National Health Mission [Internet]. [cited 2023 Apr 25]. Available from: <https://nhm.gov.in/index1.php?lang=1&level=2&sublinkid=822&lid=218>
25. Pradhan Mantri Surakshit Matritva Abhiyan | PMSMA [Internet]. [cited 2023 Apr 25]. Available from: <https://pmsma.mohfw.gov.in/>
26. Munn Z, Moola S, Lisy K, Riitano D, Tufanaru C. Chapter 5: Systematic reviews of prevalence and incidence. In: Aromataris E, Munn Z (Editors). *JBIM Manual for Evidence Synthesis*. JBI, 2020 [Internet]. [cited 2023 Jun 23]. Available from: <https://jbi.global/critical-appraisal-tools>

27. National Family Health Survey [Internet]. [cited 2023 Jun 23]. Available from: <http://rchiips.org/nfhs/>
28. Mukhopadhyay D, Bhattacharjee S, Mukhopadhyay S, Malik S, Nayak S, Biswas A. Birth preparedness and complication readiness among women of Bankura District, West Bengal. *J Fam Med Prim Care* [Internet]. 2016 [cited 2023 Mar 24];5(2):404. Available from: <https://journals.lww.com/10.4103/2249-4863.192352>
29. Girma D, Walelign A, Dejene H. Birth preparedness and complication readiness practice and associated factors among pregnant women in Central Ethiopia, 2021: A cross-sectional study. *PLOS ONE* [Internet]. 2022 Oct 27 [cited 2023 Jun 14];17(10):e0276496. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC9612452/>
30. Margaret O. Akinwaare, Abimbola Oluwatosin, Olakekan Uthman, Elizabeth Ike. WOMEN'S BIRTH PREPAREDNESS AND COMPLICATION READINESS IN NIGERIA: A SYSTEMATIC REVIEW AND META-ANALYSIS. *medRxiv* [Internet]. 2021 Jan 1;2021.11.03.21265854. Available from: <http://medrxiv.org/content/early/2021/11/03/2021.11.03.21265854.abstract>
31. Gedefa AG, Bekele AA, Kitila KM, Eba LB. Barriers to birth preparedness and complication readiness among pregnant women in rural Ethiopia: using a mixed study design, 2020. *BMJ Open* [Internet]. 2023 Apr 1 [cited 2023 Jun 14];13(4):e069565. Available from: <https://bmjopen.bmj.com/content/13/4/e069565>
32. Nimavat KA, Mangal AD, Unadkat SV, Yadav SB. A study of birth preparedness of women in Jamnagar district, Gujarat, India. *Int J Community Med Public Health* [Internet]. 2016 Dec 24 [cited 2023 Jun 15];3(9):2403–8. Available from: <https://www.ijcmph.com/index.php/ijcmph/article/view/182>

33. Berhe AK, Muche AA, Fekadu GA, Kassa GM. Birth preparedness and complication readiness among pregnant women in Ethiopia: a systematic review and Meta-analysis. *Reprod Health* [Internet]. 2018 Oct 29 [cited 2023 Jun 14];15(1):182. Available from: <https://doi.org/10.1186/s12978-018-0624-2>
34. Moran AC, Sangli G, Dineen R, Rawlins B, Yaméogo M, Baya B. Birth-Preparedness for Maternal Health: Findings from Koupéla District, Burkina Faso. *J Health Popul Nutr* [Internet]. 2006 Dec [cited 2023 Jun 15];24(4):489–97. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3001153/>

Dissertation

ORIGINALITY REPORT

| | | | |
|------------------|------------------|--------------|----------------|
| 11 % | 10 % | 8 % | 5 % |
| SIMILARITY INDEX | INTERNET SOURCES | PUBLICATIONS | STUDENT PAPERS |

PRIMARY SOURCES

| | | |
|----------|---|----------------|
| 1 | www.ncbi.nlm.nih.gov Internet Source | 2 % |
| 2 | www.medrxiv.org Internet Source | 1 % |
| 3 | oup.silverchair-cdn.com Internet Source | 1 % |
| 4 | jisponline.com Internet Source | 1 % |
| 5 | reproductive-health-journal.biomedcentral.com Internet Source | 1 % |
| 6 | Submitted to <u>University</u> of Chichester Student Paper | 1 % |
| 7 | Submitted to <u>University</u> of Melbourne Student Paper | 1 % |
| 8 | jfmipc.com Internet Source | <1 % |
| 9 | pure.rug.nl Internet Source | <1 % |

| | | |
|----|--|-----|
| 10 | Submitted to <u>University of Birmingham</u> Student Paper | <1% |
| 11 | www.frontiersin.org Internet Source | <1% |
| 12 | Submitted to CVC Nigeria Consortium Student Paper | <1% |
| 13 | Submitted to London School of Hygiene and Tropical Medicine Student Paper | <1% |
| 14 | onlinelibrary.wiley.com Internet Source | <1% |
| 15 | serval.unil.ch Internet Source | <1% |
| 16 | ijhas.in Internet Source | <1% |
| 17 | www.slideshare.net Internet Source | <1% |
| 18 | Yunga Patience Ijang, Samuel Nambile Nambile Cumber, Claude Ngwayu Nkfusai, Mbinkar Adeline Venyuy et al. "Awareness and practice of birth preparedness and complication readiness among pregnant women in the Bamenda Health District, Cameroon", BMC Pregnancy and Childbirth, 2019 Publication | <1% |

| | | |
|----|---|-----|
| 19 | dx.doi.org Internet Source | <1% |
| 20 | etheses.bham.ac.uk Internet Source | <1% |
| 21 | www.emeraldinsight.com Internet Source | <1% |
| 22 | www.panafrican-med-journal.com Internet Source | <1% |
| 23 | www.researchgate.net Internet Source | <1% |
| 24 | "RSSDI Clinical Practice Recommendations for the Management of Type 2 Diabetes Mellitus 2022", International Journal of Diabetes in Developing Countries, 2022 Publication | <1% |
| 25 | Margaret Omowaleola Akinwaare, Abimbola Oluwatosin, Olakekan Uthman, Elizabeth Ike. "Women's Birth Preparedness and Complication Readiness in Nigeria: A Systematic Review and Meta-Analysis", International Journal of Childbirth, 2023 Publication | <1% |
| 26 | journals.innovareacademics.in Internet Source | <1% |

ANNEXURE

Birth preparedness and complication readiness among pregnant women in India: Systematic review and Meta-analysis

To enable PROSPERO to focus on COVID-19 submissions, this registration record has undergone basic automated checks for eligibility and is published exactly as submitted. PROSPERO has never provided peer review, and usual checking by the PROSPERO team does not endorse content. Therefore, automatically published records should be treated as any other PROSPERO registration. Further detail is provided [here](#).

Citation

Tanya Singh, Brajaraj Tripathy, Anuj Kumar Pandey, Diksha Gautam, Sidharth Sekhar Mishra, Sutapa Bandyopadhyay Neogi. Birth preparedness and complication readiness among pregnant women in India: Systematic review and Meta-analysis. PROSPERO 2023 CRD42023396109 Available from:

https://www.crd.york.ac.uk/prospero/display_record.php?ID=CRD42023396109

Review question

Does Birth preparedness and complication readiness interventions have good birth outcome in India?

As a Mandate For SRMA this was registered on PROSPERO - Registration No. is – CRD42023396109

Keywords

| | |
|---|---|
| Title: Birth preparedness and complication readiness among pregnant women in India: systematic review and Meta-analysis | |
| Inclusion Criteria | Cross-sectional studies |
| Population | Pregnant women: "pregnan*"[Title/Abstract] OR "pregnant women"[Title/Abstract] OR "antenatal"[Title/Abstract] OR "anc"[Title/Abstract] |
| Intervention | - |
| Comparison | - |
| Outcome | Birth preparedness: "birth preparedness"[Title/Abstract] OR "preparedness"[Title/Abstract] OR "preparing for birth"[Title/Abstract] OR "emergency preparedness"[Title/Abstract] OR "birth plan"[Title/Abstract] |
| | Complication Readiness: "danger signs"[Title/Abstract] OR "readiness"[Title/Abstract] OR ("recognisable"[All Fields] OR "recognise"[All Fields] OR "recognised"[All Fields] OR "recognises"[All Fields] OR "recognising"[All Fields] OR "recognize"[All Fields] OR "recognized"[All Fields] OR "recognizes"[All Fields] OR "recognizing"[All Fields]) AND "danger sign"[Title/Abstract]) OR "obstetric complication"[Title/Abstract] OR "pregnancy complication"[Title/Abstract] OR "obstetric danger sign"[Title/Abstract] OR "maternal complications"[Title/Abstract] OR "maternal health"[Title/Abstract] OR "newborn health"[Title/Abstract] |

| Title | | | |
|---------------------------------------|-----------|---|----------------|
| Database | No | Search Query | Results |
| PUBMED (Data as on 06-02-2023) | | | |
| | #1 | "pregnan*"[Title/Abstract] OR "pregnant women"[Title/Abstract] OR "antenatal"[Title/Abstract] OR "anc"[Title/Abstract] | 620,091 |
| | #2 | "birth preparedness"[Title/Abstract] OR "preparedness"[Title/Abstract] OR "preparing for birth"[Title/Abstract] OR "emergency preparedness"[Title/Abstract] OR "birth plan"[Title/Abstract] | 20,978 |
| | #3 | "danger signs"[Title/Abstract] OR "readiness"[Title/Abstract] OR (("recognisable"[All Fields] OR "recognise"[All Fields] OR "recognised"[All Fields] OR "recognises"[All Fields] OR "recognising"[All Fields] OR "recognize"[All Fields] OR "recognized"[All Fields] OR "recognizes"[All Fields] OR "recognizing"[All Fields]) AND "danger sign"[Title/Abstract]) OR "obstetric complication"[Title/Abstract] OR "pregnancy complication"[Title/Abstract] OR "obstetric danger sign"[Title/Abstract] OR "maternal complications"[Title/Abstract] OR "maternal health"[Title/Abstract] OR "newborn health"[Title/Abstract] | 39,433 |
| | #4 | #1 AND #2 AND #3 | 276 |

| Title | | | |
|---|-----------|---|----------------|
| Database | No | Search Query | Results |
| PROQUEST (Data as on 06-02-2023) | | | |
| | #1 | noft(pregnan*) OR noft(pregnant women) OR noft(antenatal) OR noft(anc) | 124,342 |
| | #2 | noft(birth preparedness) OR noft(preparedness) OR noft(preparing for birth) OR noft(emergency preparedness) OR noft(birth plan) | 29,912 |
| | #3 | noft(complication readiness) OR noft(readiness) OR noft(recognising danger signs) OR noft(danger sign) OR noft(obstetric complication) OR noft(pregnancy complication) OR noft(obstetric danger sign) OR noft(maternal complications) OR noft(maternal health) OR noft(newborn health) | 95,890 |
| | #4 | #1 AND #2 AND #3 (noft(pregnan*) OR noft(pregnant women) OR noft(antenatal) OR noft(anc)) AND (noft(birth preparedness) OR noft(preparedness) OR noft(preparing for birth) OR noft(emergency preparedness) OR noft(birth plan)) AND (noft(complication readiness) OR noft(readiness) OR noft(recognising danger signs) OR noft(danger sign) OR noft(obstetric complication) OR noft(pregnancy complication) OR noft(obstetric danger sign) OR noft(maternal complications) OR noft(maternal health) OR noft(newborn health)) | 848 |

| Title | | | |
|---|-----------|--|--|
| Database | No | Search Query | Results |
| Cochrane (Data as on 06-02-2023) | | | |
| | #1 | (pregnan*):ti,ab OR (pregnant women):ti,ab OR (antenatal):ti,ab OR (anc):ti,ab | 68406 |
| | #2 | (birth preparedness):ti,ab OR (preparedness):ti,ab OR (preparing for birth):ti,ab OR (emergency preparedness):ti,ab OR (birth plan):ti,ab | 1639 |
| | #3 | (complication readiness):ti,ab OR (readiness):ti,ab OR (recognising danger signs):ti,ab OR (danger sign):ti,ab OR (obstetric complication):ti,ab OR (pregnancy complication):ti,ab OR (obstetric danger sign):ti,ab OR (maternal complications):ti,ab OR (maternal health):ti,ab OR (newborn health):ti,ab | 14641 |
| | #4 | #1 AND #2 AND #3 | 225 (193 - TRIALS, COCHRANE REVIEWS - 32) |

Title and Abstract Screening using RAYAAN.AI Software

longitudinal 29

randomised controlled trials 27

observational 27

reviews 27

More >>

Labels

SRMA 19

location 7

study design 4

summary read 3

no full text 2

preg women ?? 1

followup study 1

Exclusion reasons

wrong title 486

wrong population 396

wrong study design 69

foreign language 7

Topics

Nepal Bangladesh Kenya

Humans Maternal Welfare

Nigeria Only Child Child Infant

Cesarean Section

Pregnancy

2023-02-07: Birth preparedness and complication readiness among pregnant women in India: systematic review and Meta-analysis Blind ON

Detect duplicates Compute ratings Export Copy New search All reviews

Showing 1 to 6 of 971 unique entries Search: id or title or abstract or author

| Date | | Title | Authors | Rating |
|------------|--|--|---------------------------------|--------|
| 2015-01-01 | Brajara wrong study design | Health system and community level interventions for improving ... | Mbuagbaw, L; Medley, N; Da... | |
| 2020-01-01 | Brajara wrong study design | Targeted client communication via mobile devices for improving ... | Palmer, MJ; Henschke, N; Be... | |
| 2020-01-01 | Brajara wrong study design | Strategies for optimising antenatal corticosteroid administration... | Rohwer, AC; Oladapo, OT; H... | |
| 2007-01-01 | Brajara wrong study design | Individual or group antenatal education for childbirth or parenth... | Gagnon, AJ; Sandall, J | |
| 2015-01-01 | Brajara wrong study design | Group versus conventional antenatal care for women | Catling, CJ; Medley, N; Four... | |

✔ Include
? Maybe
✘ Exclude

📄 Add Note
✔ Highlights ON
📄 Upload PDF full-texts

Antenatal and Postnatal Care

Authors: Aderoba AK, Adu-Bonsaffoh K.;

Journal: - Volume 0, Issue 0, pp. - published

Publication Types: Journal Article

Topics: Postnatal Care

System Id: 935199820

Search methods: Uploaded References [PubMed_V1_SRMA_06022023.csv]

DOI: 10.1016/j.ogc.2022.07.005

Brajara Help

not duplicates 0

Resolved 0

Auto Deduplicated

2 exact matches 3

Inclusion decisions

Undecided 0

Maybe 0

Included 13

Excluded 958

Minimum collaborator decisions

At least 1 971

At least 2 971

At least 3 0

Maximum collaborator decisions

At most 0 0

At most 1 0

At most 2 971

Search methods [Add new]

Uploaded References [final co... 32

Uploaded References [ProQue... 848

Uploaded References [PubMed... 276

Keywords for include [Add new]

pregnant women 262

Birth Preparedness 219

cross-sectional 142

Complication Readiness 131

Amnion etc 60

2023-02-07: Birth preparedness and complication readiness among pregnant women in India: systematic review and Meta-analysis Blind ON

Detect duplicates Compute ratings Export Copy New search All reviews

Showing 1 to 6 of 971 unique entries Search: id or title or abstract or author

| Date | | Title | Authors | Rating |
|------------|--|--|---------------------------------|--------|
| 2015-01-01 | Brajara wrong study design | Health system and community level interventions for improving ... | Mbuagbaw, L; Medley, N; Da... | |
| 2020-01-01 | Brajara wrong study design | Targeted client communication via mobile devices for improving ... | Palmer, MJ; Henschke, N; Be... | |
| 2020-01-01 | Brajara wrong study design | Strategies for optimising antenatal corticosteroid administration... | Rohwer, AC; Oladapo, OT; H... | |
| 2007-01-01 | Brajara wrong study design | Individual or group antenatal education for childbirth or parenth... | Gagnon, AJ; Sandall, J | |
| 2015-01-01 | Brajara wrong study design | Group versus conventional antenatal care for women | Catling, CJ; Medley, N; Four... | |

✔ Include
? Maybe
✘ Exclude

📄 Add Note
✔ Highlights ON
📄 Upload PDF full-texts

Antenatal and Postnatal Care

Authors: Aderoba AK, Adu-Bonsaffoh K.;

Journal: - Volume 0, Issue 0, pp. - published

Publication Types: Journal Article

Topics: Postnatal Care

System Id: 935199820

Search methods: Uploaded References [PubMed_V1_SRMA_06022023.csv]

DOI: 10.1016/j.ogc.2022.07.005

Brajara Help

Possible Duplicates

| | |
|----------------|-----|
| Unresolved | 0 |
| Deleted | 182 |
| Not duplicates | 3 |
| Resolved | 4 |

Auto Deduplicated

2 exact matches

Inclusion decisions [Clear]

| | |
|-----------|-----|
| Undecided | 0 |
| Maybe | 6 |
| Included | 13 |
| Excluded | 949 |
| Conflict | 3 |

Decision by

Doctor Sidharth Mishra
Brajaraaj
Tanya

Minimum collaborator decisions

| | |
|------------|-----|
| At least 1 | 971 |
| At least 2 | 965 |
| At least 3 | 0 |

Maximum collaborator decisions

| | |
|-----------|-----|
| At most 0 | 0 |
| At most 1 | 6 |
| At most 2 | 971 |

2023-02-07: Birth preparedness and complication readiness among pregnant women in India: systematic review and Meta-analysis Blind OFF

Detect duplicates Compute ratings Export Copy New search All reviews

Showing 1 to 6 of 13 unique entries (filtered from 971 total unique entries) Search:

| Date | Title | Authors | Rating |
|--|--|--------------------------------|--------|
| Brajaraaj Tanya cross sectional study | Birth preparedness and complication readiness among wom... | Mukhopadhyay DK, Bhattach... | |
| Brajaraaj Tanya | Birth preparedness and complication readiness in pregnant women attending... | Kamineni V, Murki AD, Kota ... | |
| Brajaraaj Tanya study design | Status and determinants of birth preparedness and complication r... | Sharma N, Kumar N, Singh ... | |
| Brajaraaj Tanya | Status of birth preparedness and complication readiness in Uttar Dinajpur Dis... | Mukhopadhyay DK, Mukhop... | |
| Brajaraaj Tanya | Status of birth preparedness & complication readiness in Rewa District of Mad... | Kushwah SS, Dubey D, Sing... | |

No articles selected, use your mouse or keyboard to select articles from the above table.

Help

REVIEW CHAT

66 | Page

CHECKLIST FOR RISK OF BIAS

JBI CRITICAL APPRAISAL CHECKLIST FOR STUDIES REPORTING PREVALENCE DATA

| | Yes | No | Unclear | Not applicable |
|---|--------------------------|--------------------------|--------------------------|--------------------------|
| 1. Was the sample frame appropriate to address the target population? | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| 2. Were study participants sampled in an appropriate way? | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| 3. Was the sample size adequate? | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| 4. Were the study subjects and the setting described in detail? | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| 5. Was the data analysis conducted with sufficient coverage of the identified sample? | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| 6. Were valid methods used for the identification of the condition? | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| 7. Was the condition measured in a standard, reliable way for all participants? | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |